

9. THE ALGEBRA-GEOMETRY DICTIONARY

Goal: study the correspondence

$$\begin{array}{ccc} \text{varieties} & & \text{ideals} \\ W & \longrightarrow & I(W) \\ & & \{p : p(a) = 0 \quad \forall a \in W\} \end{array}$$

$$\begin{array}{ccc} V(J) & \longleftarrow & J \\ \{a : p(a) = 0 \quad \forall p \in J\} & & \end{array}$$

Example 9.1

The correspondence is not a bijection:

(1) For $I = \langle x \rangle$, $J = \langle x^m \rangle$ in $k[x]$
 $V(I) = \{0\} = V(J)$

More generally for any $p \in k[x_1, \dots, x_n]$ and $m \in \mathbb{N}$
 $V(p) = V(p^m)$ but $\langle p^m \rangle \subsetneq \langle p \rangle$

(2) Non-algebraically closed fields are more problematic!
In $\mathbb{R}[x]$ for $I = \langle 1+x^2 \rangle$, $J = \langle 1+x^2+x^4 \rangle$
 $I \not\subseteq J$, $J \not\subseteq I$, $V(I) = V(J) = \emptyset$

Similarly in $\mathbb{R}[x, y]$

$$\begin{aligned} V(1+x^2+y^2) &= V(1+x^2+y^4) = V(1+x^2y^2) = \emptyset \\ \langle 1+x^2+y^2 \rangle &\neq \langle 1+x^2+y^4 \rangle \neq \langle 1+x^2y^2 \rangle \end{aligned}$$

(3) Suppose $I \subset K[x]$ with $V(I) = \emptyset$.

Univariate ideals are principal, so

$$I = \langle p \rangle, \quad p \in K[x] \Rightarrow V(p) = V(I) = \emptyset$$

If K algebraically closed, then

$$\begin{aligned} V(p) = \emptyset &\Rightarrow p \text{ nonzero constant} \\ &\Rightarrow I = \langle 1 \rangle = K[x] \end{aligned}$$

Theorem 9.2 (Weak Nullstellensatz)

Let K be algebraically closed

and $I \subset K[x_1, \dots, x_n]$ an ideal. Then

$$V(I) = \emptyset \Leftrightarrow I = K[x_1, \dots, x_n]$$

Proof

" \Leftarrow " is immediate.

The nontrivial claim is $I \subsetneq K[x_1, \dots, x_n] \Rightarrow V(I) \neq \emptyset$.

This follows by induction on $n \geq 1$.

The case $n=1$ is Example 9.1(3).

For the induction step we will prove

$$\textcircled{*} \quad I \subsetneq K[x_1, \dots, x_n] \Rightarrow \exists a \in K \text{ s.t. } I_{x_n=a} \subsetneq K[x_1, \dots, x_{n-1}]$$

where

$$I_{x_n=a} = \{ p(x_1, \dots, x_{n-1}, a) \in K[x_1, \dots, x_n] : p \in I \}.$$

The proof of $\textcircled{*}$ splits into two cases.

Case 1: $I \cap K[x_n] \neq \{0\}$.

Let $0 \neq p \in I \cap K[x_n]$.

Since K is algebraically closed, $\exists c, a_1, \dots, a_m \in K$ st.

$$p = c \prod_{j=1}^m (x_n - a_j)$$

Note: $m \geq 1$ since otherwise $1 = \frac{1}{c} p \in I$

Claim: \textcircled{B} holds for some $c = a_j$, $j = 1, \dots, m$.

Proof: Suppose not. Then for each $j = 1, \dots, m$

$$I_{x_n=a_j} = K[x_1, \dots, x_{n-1}] \Rightarrow 1 \notin I_{x_n=a_j}$$

$$\Rightarrow \exists q_j \in I \text{ such that } q_j(x_1, \dots, x_{n-1}, a_j) = 1$$

Then $q_j = 1 + (x_n - a_j) \cdot h_j$ for some $h_j \in K[x_1, \dots, x_{n-1}]$:

$$\text{If } q_j = \sum_{\alpha} c_{\alpha} x^{\alpha} x_n^{\alpha_n}, \quad \alpha = (\alpha_1, \dots, \alpha_{n-1}, \alpha_n) = (\bar{\alpha}, \alpha_n)$$

$$\begin{aligned} \text{then } q_j(x_1, \dots, x_{n-1}, a_j + x_n a_j) &= \sum_{\alpha} c_{\alpha} \bar{x}^{\alpha} (a_j + (x_n - a_j))^{\alpha_n} \\ &= \sum_{\alpha} c_{\alpha} \left(\bar{x}^{\alpha} a_j^{\alpha_n} + (x_n - a_j)(\dots) \right) \end{aligned}$$

It follows that

$$\begin{aligned} 1 &= \prod_{j=1}^m (q_j - (x_n - a_j) \cdot h_j) \in \prod_{j=1}^m (I + (x_n - a_j) h_j) \\ &\subset \prod_{j=1}^m (x_n - a_j) h_j + I \\ &= \frac{1}{c} p \cdot \prod_{j=1}^m h_j + I \subset I. \quad \checkmark \end{aligned}$$

Hence \textcircled{B} must hold for some $j = 1, \dots, m$

Case 2: $I \cap K[x_n] = 0$

Let $G = \{g_1, \dots, g_t\}$ Gröbner basis of I in lex.

Decompose the leading monomials as

$$LM(g_i) = x^{\alpha_i} \cdot x_n^{\beta_i}, \quad x^{\alpha_i} \text{ monomial in } x_1, \dots, x_{n-1}$$

and collect all terms with a x^{α_i} factor

(*) $g_i = c_i(x_n) \cdot x^{\alpha_i} + \dots \text{ terms } < x^{\alpha_i}$
where $0 \neq c_i \in K[x_n]$

Let $a \in K$ be such that $c_i(a) \neq 0 \quad \forall i=1, \dots, t$

(Note: this exists since an algebraically closed field is infinite)

Since G is a basis of I ,

$$\bar{g}_i := g_i(x_1, \dots, x_{n-1}, a) \in K[x_1, \dots, x_{n-1}], \quad i=1, \dots, t$$

is a basis of $\overline{I}_{x_n=a}$.

Claim: $\bar{G} = \{\bar{g}_1, \dots, \bar{g}_t\}$ is a Gröbner basis of $I_{x_n=a}$.

Proof: We will use the lcm -representation generalization of Buchberger's criterion.

For $1 \leq i, j \leq t$, let $x^\gamma = lcm(x^{\alpha_i}, x^{\alpha_j})$ and consider

$$S = c_j(x_n) \cdot \frac{x^\gamma}{x^{\alpha_i}} g_i - c_i(x_n) \frac{x^\gamma}{x^{\alpha_j}} g_j$$

By (*) $LT(S) < x^\gamma$.

By polynomial division we get a standard representation

$$S = \sum_{l=1}^t q_l g_l, \quad LT(q_l g_l) \leq LT(S)$$

Evaluating at $x_n = a$, we get

$$\bar{S} = C_j(a) \frac{x^\gamma}{x^{\alpha_i}} \bar{g}_i - C_i(a) \frac{x^\gamma}{x^{\alpha_j}} \bar{g}_j = \sum_{l=1}^t \bar{q}_l \bar{g}_l$$

$$\text{where } \bar{q}_l = q_l(x_1, \dots, x_{n-1}, a) \in k[x_1, \dots, x_{n-1}]$$

Hence we have

- $\bar{S} = C_i(a)C_j(a) \cdot S(\bar{g}_i, \bar{g}_j)$
- $LT(\bar{q}_l \bar{g}_l) \leq LT(q_l g_l) \leq LT(S) < x^\gamma$
- $x^\gamma = \text{lcm}(x^{\alpha_i}, x^{\alpha_j}) = \text{lcm}(LM(\bar{g}_i), LM(\bar{g}_j))$

So each S -polynomial $S(\bar{g}_i, \bar{g}_j) \in k[x_1, \dots, x_{n-1}]$
has a lcm-representation $\Rightarrow \bar{G}$ Gröbner basis.

To conclude the proof, observe that

$$LT(\bar{g}_i) = C_i(a) x^{\alpha_i}$$

is not a constant since otherwise

$$LM(g_i) = x^{\alpha_i} x_n^{m_i} = x_n^{m_i} \Rightarrow g_i \in k[x_n]$$

and then $I \cap k[x_n] = 0 \Rightarrow g_i = 0$.

Hence $LT(\bar{g}_i) \neq 1$ for all $i \neq 1, \dots, t$

so $1 \notin I|_{x_n=a}$. \square

Fundamental Theorem of algebra:

$$p \in \mathbb{C}[t] \quad \& \quad 1 \notin \langle p \rangle \Rightarrow \exists \text{ solution to } p=0$$

Weak Nullstellensatz:

$$\begin{aligned} p_1, \dots, p_m &\in \mathbb{C}[x_1, \dots, x_n] \quad \& \quad 1 \notin \langle p_1, \dots, p_m \rangle \\ \Rightarrow \exists \text{ solution to } p_1 = \dots = p_m = 0 \end{aligned}$$

Theorem 9.3 (Hilbert's Nullstellensatz)

Let K be an algebraically closed field

and $I = \langle p_1, \dots, p_s \rangle \subset K[x_1, \dots, x_n]$. Then

$$f \in I(V(I)) \iff f^m \in I \text{ for some } m \in \mathbb{N}$$

Proof

" \Leftarrow " If $f^m \in I$, then $f^m(c) = 0 \quad \forall c \in V(I)$

so also $f(c) = 0 \quad \forall c \in V(I)$.

" \Rightarrow " Rabinowitsch's trick: Consider the ideal

$$J := \langle p_1, \dots, p_s, 1-yf \rangle \subset K[x_1, \dots, x_n, y]$$

Claim: $V(J) = \emptyset$

Prof: Let $(a, b) \in K^n \times K$. Either $a \in V(I)$ or $a \notin V(I)$.

If $a \in V(I)$ then by assumption $f(a) = 0$, so

$$(1-yf)(a, b) = 1 - b \cdot f(a) = 1 \neq 0 \Rightarrow (a, b) \notin V(J)$$

If $a \notin V(I)$ then $p_i(a, b) = p_i(a) \neq 0$ for some $i = 1, \dots, s$

so again $(a, b) \in V(J)$.

Apply the weak Nullstellensatz to obtain $l \in J$, so

$$1 = \sum_{i=1}^s q_i p_i + q(1-yf)$$

for some $q_1, \dots, q_n, q \in K[x_1, \dots, x_n, y]$

Formally substituting $y = 1/f(x_1, \dots, x_n)$ we get
a rational expression in x_1, \dots, x_n

$$1 = \sum_{i=1}^s q_i(x_1, \dots, x_n, \frac{1}{f}) p_i(x_1, \dots, x_n)$$

Clearing denominators, we obtain a polynomial identity

$$f^m = \sum_{i=1}^s \tilde{q}_i p_i \Rightarrow f^m \in I. \quad \square$$

$I(V)$ is a special type of ideal:

$$f^m \in I(V) \Rightarrow f \in I(V)$$

Definition 9.4

(1) An ideal I is radical if $f^n \in I \Rightarrow f \in I$.

(2) The radical of an ideal I is the set

$$\sqrt{I} = \{f : f^m \in I \text{ for some } m \in \mathbb{N}\}$$

Example 9.5

Let $I = \langle x^2, y^3 \rangle \subset k[x, y]$, so

$$x \in \sqrt{I} \quad \text{and} \quad y \in \sqrt{I}$$

Then also

$$xy \in \sqrt{I} \quad \text{since} \quad (xy)^2 = x^2 \cdot y^2 \in I$$

and by the binomial formula $x+y \in \sqrt{I}$ since

$$(x+y)^4 = \underbrace{x^4}_{\substack{\text{multiples of } x^2}} + 4x^3y + 6x^2y^2 + 4xy^3 + y^4 \in I \quad \begin{matrix} \nearrow \\ \text{multiples of } y^3 \end{matrix}$$

Lemma 9.6

If $I \subset k[x_1, \dots, x_n]$ is an ideal,

then \sqrt{I} is a radical ideal with $I \subset \sqrt{I}$.

Proof

$I \subset \sqrt{I}$ is immediate (take $m=1$)

That \sqrt{I} is an ideal follows by the argument
of Example 9.5:

$$p \in \sqrt{I} \Rightarrow p^m \in I \Rightarrow (pq)^m \in I \quad \text{for all } q$$

$$\begin{aligned} p, q \in \sqrt{I} &\Rightarrow p^m \in I \quad \text{and} \quad q^l \in I \\ &\Rightarrow (p+q)^{m+l-1} = \sum_{i=0}^{m+l-1} \binom{m+l-1}{i} p^i q^{m+l-1-i} \in I \end{aligned}$$

Since each summand has either $i \geq m$ or
 $i \leq m-1 \Rightarrow m+l-1-i \geq l$.

Finally, $f^m \in \sqrt{I} \Rightarrow f^{ml} \in I \Rightarrow f \in \sqrt{I} \quad \text{so } \sqrt{I} \text{ is radical.} \square$

Theorem 9.7 (Strong Nullstellensatz)

Let k be algebraically closed

and $I \subset k[x_1, \dots, x_n]$ an ideal. Then

$$I(V(I)) = \sqrt{I}$$